arXiv:1411.3323v2 [cond-mat.str-el] 26 Jul 2016

Charge density waves in graphite; towards the magnetic ultra-quantum limit

F. Arnold,[1,*] A. Isidori,[1] E. Kampert,[2] B. Yager,[1] M. Eschrig,[1] and J. Saunders[1]

[1] *Royal Holloway, University of London, TW20 0EX Egham, United Kingdom*
[2] *Hochfeld-Magnetlabor Dresden (HLD), Helmholtz-Zentrum Dresden-Rossendorf, D-01328 Dresden, Germany*

(Dated: July 27, 2016)

Graphite is a model system for the study of three-dimensional electrons and holes in the magnetic quantum limit, in which the charges are confined to the lowest Landau levels. We report magneto-transport measurements in pulsed magnetic fields up to 60 T, which resolve the collapse of two density wave states in two, electron and hole, Landau levels at 52.3 and 54.2 T respectively. We report evidence for a commensurate density wave at 47.1 T in the electron Landau level. The theoretical modelling of these results predicts that the ultra-quantum limit is entered above 73.5 T. This state is an insulator, and we discuss its correspondence to the "metallic" state reported earlier. We propose that this (interaction-induced) insulating phase supports surface states that carry no charge or spin within the planes, but does however support charge transport out of plane.

Semimetals like graphite and bismuth are the subject of renewed interest due to their close relation to the topological Dirac and Weyl semimetals [1–3]. Their low charge carrier density and effective mass reduce the magnetic field necessary to drive these systems into their magnetic quantum limit, in which the electronic structure is described by only a few Landau levels (LLs). In such materials this limit can be realised in state-of-the-art high-magnetic field laboratories, whereas it is inaccessible in conventional metals. The magnetic field induces a crossover from 3D to 1D physics, and the perfect nesting characteristic of 1D electron systems leads to the possibility of field-induced spin- and charge-density wave instabilities (SDWs and CDWs, respectively) with ordering wave-vector along the field direction [4–7]. In the ultra-quantum regime these systems are potentially subject to a rich variety of magnetic field driven topological quantum states such as fractional quantum Hall effect and the recently discussed axial and chiral anomaly [7–13]. The clarification of the nature of the density wave instabilities in graphite, and the nature of the quantum state at high magnetic fields, is the focus of the experimental and theoretical work reported here.

Graphite is a well known 3D semimetal consisting of an infinite stack of graphene sheets [14, 15]. The zero-field band structure is well described by the tight-binding Slonczewski-Weiss-McClure model [16–19]. The Fermi surface consists of strongly anisotropic, trigonally warped ellipsoidal electron and hole pockets located along the H-K-H and H'-K'-H' edges of the hexagonal Brillouin zone (see Fig. 1). In magnetic fields applied perpendicular to the graphene layers, the E_3-band becomes slightly hybridised with the E_1 and E_2 bands in proximity of the H points and gives rise to two 1D LLs with index $n = -1$ and $n = 0$. Each LL is spin-split by the Zeeman energy and has a residual twofold valley-degeneracy associated with the two inequivalent H-K-H and H'-K'-H' edges. For magnetic fields above 8 T these are the only LLs that cross the Fermi energy. The system becomes quasi-1D, due to c-axis dispersion.

This quantum limit has been extensively investigated by magneto-transport measurements. The first studies by Iye *et al.* on Kish graphite showed a pronounced resistance anomaly above 22 T [20]. Nakao [21], Yoshioka and Fukuyama [22] attributed the observed anomaly to the formation of a CDW in one of the lower LLs and developed a mean-field theory describing the magnetic field dependence of its critical temperature. Improved LL calculations by Takada and Goto [23] showed that electron correlations play a crucial role in renormalizing the bare (free-electron) Slonczewski-Weiss-McClure LL structure. Subsequent experiments to higher fields [24, 25] identified a transition at 52 T corresponding to the collapse of a density wave phase as the $(0,\uparrow)$ or $(-1,\downarrow)$ LL emptied. So far attempts to directly measure the gap of this putative density wave spectroscopically [26, 27] have been unsuccessful.

Recently, Fauqué et al. [28] discovered the onset of a second density-wave anomaly above 53 T, by out-of-plane transport measurements on Kish graphite up to 80 T. This state was found to collapse at 75 T into a state with metallic conductivity. These authors argued that only one of the $(0,\uparrow)$ or $(-1,\downarrow)$ LLs depopulates at 53 T, and furthermore that the $(0,\downarrow)$ and $(-1,\uparrow)$ levels



Figure 1. Zero-field k_z-dependence of graphite's band-structure along the H-K-H and H'-K'-H' edges of the Brillouin zone.



Figure 2. a) Magnetic field dependence of the in-plane resistance of graphite for temperatures below 10 K measured in a 25 ms 60 T pulsed field magnet. b) Magnetic field derivatives of the in-plane resistance: The data has been offset proportional to the temperature by 0.5 Ω/TK for clarity. α and β denote the onset transitions of the CDWs in the $(0,\uparrow)$ and $(-1,\downarrow)$, whereas α' and β' mark the field at which they vanish. At γ the CDW associated with the $(0,\uparrow)$ LL undergoes a lock-in transition.

remain populated above 75 T. On the other hand, the observation of a vanishing Hall coefficient at 53 T [29] was interpreted in terms of the depopulation of both the $(0,\uparrow)$ and $(-1,\downarrow)$ LLs at 53 T, as predicted theoretically [23]. It was suggested that an excitonic phase forms in the remaining $(0,\downarrow)$ and $(-1,\uparrow)$ levels above 53 T . Elsewhere it has been argued that an excitonic insulator phase appears at 46 T [30]. It is clear that the evolution of the LLs with magnetic field, and their associated density wave instabilities, through the quasi-quantum limit and into the ultra-quantum limit at highest fields, remain open questions.

In this article we present new magneto-transport measurements on single crystal graphite, of significantly higher quality than hitherto, in conjunction with new theoretical calculations of the LL band structure, and density wave instabilities. We show that the field-induced density wave below 54.2 T, contrary to previous reports,



Figure 3. Phase diagram of the CDWs in the $(0,\uparrow)$ and $(-1,\downarrow)$ LLs. Circles and squares are phase transitions with positive and negative differential magnetoresistance, where black represents the α, blue the β, and ocher the γ-transition. At low fields the critical temperature of α and β increases exponentially following Yoshioka-Fukuyama's theory [22] (black and blue dashed lines): $T_c(B) = T^* \exp\{-B^*/B\}$, with $T^*_\alpha = 230$ K, $B^*_\alpha = 148$ T, $T^*_\beta = 300$ K, $B^*_\beta = 195$ T.

is a superposition state of two incommensurate collinear DWs with different onset and vanishing fields. One of these undergoes a lock-in transition at 47.1 T before both DWs vanish due to the emptying of their corresponding LLs at 52.3 and 54.2 T respectively. These results allow us to predict that the threshold field for the ultra-quantum limit is 73.5 T, above which the LL structure is such that the bulk is an insulator. This corresponds to the new phase, with observed onset at 75 T, discovered in [28]. Given the observation of metallic response in that phase [28], we propose that it supports a set of surface states which allow for charge transport perpendicular to the planes, although within the planes the charge- and spin currents cancel. We find that interactions are crucial for this state to be insulating; without taking into account interactions the system would be metallic at these magnetic fields.

Our electrical transport measurements were performed on a single crystal of Tanzanian natural graphite [31], the quality of which significantly exceeds the commonly used Kish graphite, highly-oriented-pyrolytic graphite, or other natural graphites. We believe this sample quality is crucial in revealing the new features we observe; the self-doping of graphite by structural defects is minimised, reducing electron-hole charge imbalance. Our studies of the in-plane and out-of-plane magnetotransport were made in pulsed magnetic fields up to 60 T [32, 33].

Figure 2 shows the in-plane resistance of the graphite crystal for fields up to 60 T and temperatures below 10 K. The in-plane resistance first increases steeply, superimposed by Shubnikov-de Haas oscillations, and saturates above 15 T due to the presence of a closed Fermi surface [7, 20, 34, 35]. At lowest temperatures a step is observed in the in-plane resistance at around 30 T, followed by a steep increase of the resistance, reaching its maximum value at 48 T before it drops down to below its initial value at around 53 T. This behavior has already been reported for Kish and highly-oriented pyrolytic graphite and has been attributed to the formation of a CDW state



Figure 4. a) Renormalized LL structure in the ultra-quantum limit: Above 8 T only four LLs cross the Fermi energy. At 54 T valley degeneracy is lifted leading to two differentially doped sets of LLs, here A and B refer to the H-K-H and H'-K'-H' valleys. b) CDW nesting vectors Q up to 90 T. Hole-like levels have nesting vectors given by $Q^{(\mathrm{h})} = 2k_F^{(\mathrm{h})} = 2\pi/c_0 - 2k_F$ (see inset). All Q's decrease with increasing magnetic field due to the upward and downward shift of the electron-like $(0,\sigma)$ and hole-like $(-1,\sigma)$ LLs, before vanishing in a first-order transition at $B_{\alpha'} = 52.3$ T, $B_{\beta'} = 54.2$ T, and $B_{\delta'} = 73.5$ T. The nesting vector of the $(0,\uparrow)$ LL becomes commensurate at $B_\gamma = 48$ T. c) Close-up of the nesting vectors in the vicinity of the first-order α' and β'-transitions (top panel) and observed resistance hysteresis (bottom panel): The red curves show the magnetic field up-sweeps whereas blue curves correspond to the down-sweeps.

[20, 25, 28, 36, 37]. Similar features can also be identified in the out-of-plane transport [28, 33].

In order to highlight the signatures of the transitions in our sample, the magnetic field derivatives of the in-plane magnetoresistance were calculated (see Fig. 2). Here the transitions appear as maxima and minima. We denote the observed maxima as α, β, and γ, in ascending order of their transition fields, and the corresponding minima as α' and β'. The resolution of the two features α' and β' is the first key new observation, and is also seen in the out-of-plane transport [33]. The β' transition develops below 6 K alongside the α' transition. In addition we identify a new weaker feature, labelled γ, at 47.1 ± 0.1 T, which is resolved below 3 K and represents the onset of another resistance increase. The temperature dependence of these transitions is shown in the phase diagram of Fig. 3. [38]

We now discuss the theoretical interpretation of these features, based on our new calculations of the LL structure including the Hartree-Fock self-energy, following Takada and Goto [23]. In the quantum limit, the system is understood in terms of the one-dimensional dispersion of the lowest, spin-split, electron and hole LLs, and their density wave instabilities. We investigate closely the magnetic field dependence of the putative nesting vectors in all four LLs that are close to the Fermi level (see Fig. 3). We calculate the renormalized LLs by including the electron self-energy in a fully self-consistent manner [33]. The effect of short-range correlations, neglected by the conventional random-phase approximation (RPA), is included via the so-called local-field correction [39, 40] to the effective electron-electron interaction. The inclusion of this correction is crucial in order to get a quantitative agreement between the theoretical LL structure and the experimental data. We use the experimental value of $B_{\alpha'}$ to fix the tuning parameter that characterizes our theory [33], namely the static relative permittivity.

We find that the sharpest new features at $B_{\alpha'} = 52.3 \pm 0.1$ T and $B_{\beta'} = 54.2 \pm 0.1$ T, corresponding to two distinct first-order transitions (see Figs. 3 and 4), are associated with the abrupt depopulation of both the $(0,\uparrow)$ (electron-like) and $(-1,\downarrow)$ (hole-like) LLs, as originally predicted [23]. We believe that the DW instabilities in both LLs correspond to CDWs, and discuss fully the possible scenarios, in the light of this new direct observation of the depopulation of two LLs in [33]. We also argue below that the splitting of the transitions arises from the breaking of valley degeneracy. The first-order character of the α' and β' transitions, predicted by our theory, is experimentally confirmed by the clear evidence of hysteresis in both, as shown in Fig. 3. Note that a resistance hysteresis around 53 T was already observed by Yaguchi *et al.* [41], however the resolution of the two first-order transitions was not possible in that work.

Furthermore, our calculation finds that at $B_\gamma = 48$ T the nesting vector $Q^{(0,\uparrow)} = 2k_F^{(0,\uparrow)}$ of the $(0,\uparrow)$ LL takes the value of $2\pi/4c_0$ (see Fig. 3), corresponding to a commensurate wavelength of $4c_0$ (the lattice parameter c_0 is twice the distance between adjacent graphene layers). This suggests that the γ-feature, observed experimentally around 47.1 T and for temperatures below 3 K (see Fig. 2), corresponds to a lock-in transition of the corresponding LL from an incommensurate CDW state to a "phase-locked" commensurate order [33, 42–49], where translational invariance of the CDW modulation with respect to the lattice is lost [50]. This good agreement is achieved with no free parameter.

We now elaborate on the splitting of the α' and β' transitions. In the simplest calculation these transitions occur at exactly the same field value, in an undoped sample [51]. To account for the splitting, while preserving overall charge balance, we need to either lift the valley degener-

acy or invoke an inhomogeneous state with spatial phase separation. In the first scenario, a spontaneous breaking of the valley degeneracy leads to a splitting between the LLs originating from the Fermi surface pockets located along the two inequivalent H-K-H and H'-K'-H' edges of the Brillouin zone (see Fig. 1) similar to what has been observed in bismuth [13]. The system is then characterized by two inequivalent sets of LLs, one for each valley, allowing each set of LLs to develop a finite, equal and opposite, charge imbalance in the narrow region between $B_{\alpha'}$ and $B_{\beta'}$. In this magnetic-field region both valleys are characterized by three populated and one empty LLs; the empty level in the hole-doped valley is the electron-like $(0,\uparrow)$ LL, whereas in the electron-doped valley the empty level is the hole-like $(-1,\downarrow)$ (see Fig. 4). Since the two valleys are now characterized by inequivalent nesting vectors, the collinear CDW order observed below α' turns into a valley density-wave state, with an additional commensurate in-plane modulation originating from the reciprocal lattice vectors connecting the two valleys.

To account quantitatively for the observed splitting between α' and β' we predict a charge imbalance, parameterising the broken valley degeneracy, of $(p-n)/(p+n) = \pm 0.128$ in the two valleys, where p (n) denotes the hole (electron) carrier density. Note that the opposite imbalance in the two valleys guarantees the overall charge balance of the system.

We now turn to the nature of the system in the ultra-quantum limit. We first note that, for magnetic fields above β', the populated levels are $(0,\downarrow)$ (electron-like) and $(-1,\uparrow)$ (hole-like). These LLs are still characterized by finite nesting vectors. Note that, in the case of only two active levels, the nesting vectors arising from these bands are identical due to the charge balance between electron and hole carriers.

Our calculations, with parameters fixed by the observed field-induced transitions discussed previously, find that both levels become depopulated at 73.5 T, as shown in Fig. 4. This state was first theoretically identified by [23] in a computation at 200 T. Our predicted numerical value of $B_{\delta'}$, is in remarkable agreement with the measured critical field of 75 T for the reentrant transition of the upper density-wave phase observed by Fauqué *et al.* [28]. We note that both nesting vectors could give rise to an additional lock-in transition just below 70 T, due to their commensuration with a wavelength of $3c_0$. The state above 75 T was observed to have metallic out-of-plane conductivity [28].

In contrast with the proposal in [28], the $(0,\uparrow)$ and and $(-1,\downarrow)$ empty above 75T. In this case we expect that low-temperature transport occurs via a 2D chiral sheath of extended surface states [52–57]. These surface states are protected against weak impurity scattering. This leads to a ballistic in-plane transport along the edges of the sample, in the direction perpendicular to the magnetic field. Taking into account valley degeneracy, all four LLs contribute, leading to a cancellation of the charge and spin currents. Nevertheless metallic conductivity is observed in the out-of-plane direction [28], and we propose that this is accounted for in terms of ballistic motion in the in-plane direction, mediated by inter-layer hopping occurring at the sample edges [53–55]. At magnetic fields between 55 and 75 T the Fermi energy crosses the $(0,\uparrow)$ and $(-1,\downarrow)$ LLs. The wavefunctions of surface states have finite overlap with bulk states and are scattered by bulk impurities, which renders transport via these states diffusive, yielding a thermally activated out-of-plane conductance at temperatures below the CDW transition, as observed [28].

For even higher magnetic fields, beyond 75T, a topological phase transition into a quantum spin Hall insulating state is expected when the two bands closest to the chemical potential interchange their order due to strong Zeeman splitting. The exact field when this happens depends strongly on the type of interactions dominating the system, and cannot be predicted at this stage without further experimental input. Studying this phase transition would give insight into the interplay between strong interactions and topology in the ultra-quantum limit.

In summary, our measurements on high quality graphite samples have clarified the magnetic-phase diagram, by resolving the appearance and sudden quenching of two field induced density wave instabilities, as well as the identification of a new lock-in transition in one of these. We can theoretically account for these features and correctly predict the field-quenching of the further density wave instability observed recently [28].

It appears that in the ultra-quantum limit graphite is a fully gapped state in the bulk, an insulator, due to the strong influence of electron correlations on the LL band structure. We confirm that this state has an onset field of 75T, making it experimentally accessible in state-of-the-art pulsed magnetic field facilities. There is current interest in systems where emergent band-structure leads to a topologically non-trivial insulating state, topological Kondo insulators being one example [58, 59]. The observation of metallic out-of-plane transport [28] opens a new play ground for studying surface states in interaction-induced insulators, with the possibility of a topological phase transition at fields beyond 75T.

The authors wish to thank T. Förster, J. Wosnitza, T. Herrmansdörfer and S. Zherlitsyn for experimental support at the HLD. We acknowledge the support of the HLD-HZDR, member of the European Magnetic Field Laboratory (EMFL) as well as the Hubbard Theory Consortium and Engineering and Physical Science Research Council (EPSRC Grant Nos. EP/H048375/1 and EP/J010618/1).

* now at: Max Planck Institute for Chemical Physics of Solids, Nöthnitzer Str. 40, 01187 Dresden, Germany

[1] Z. Liu, B. Zhou, Y. Zhang, Z. Wang, H. Weng, D. Prabhakaran, S. Mo, Z. Shen, Z. Fang, X. Dai, Z. Hussain, and Y. Chen, Science **343**, 864 (2014).

[2] M. Neupane, S. Xu, R. Sankar, N. Alidoust, G. Bian, C. Liu, I. Belopolski, T. Chang, H. Jeng, H. Lin, A. Bansil, F. Chou, and M. Hassan, Nat. Comms. **5** (2014).

[3] H. Weng, C. Fang, Z. Fang, B.A. Bernevig, and X. Dia, Phys. Rev. X **5**, 011029 (2015).

[4] A. W. Overhauser, Phys. Rev. Lett. **4**, 462 (1960).

[5] A. W. Overhauser, Phys. Rev. **167**, 691 (1968).

[6] V. Celli and N. D. Mermin, Phys. Rev. **140**, A839 (1965).

[7] B. I. Halperin, Jap. J. Appl. Phys. **26**, 1913 (1987).

[8] K. Behnia, L. Balicas, and Y. Kopelevich, Science **317**, 1729 (2007).

[9] A. Banerjee, B. Fauqué, K. Izawa, A. Miyake, I. Sheikin, J. Flouquet, B. Lenoir, and K. Behnia, Phys. Rev. B **78**, 161103(R) (2008).

[10] H. Yang, B. Fauqué, L. Malone, A. Antunes, Z. Zhu, C. Uher, and K. Behnia, Nature Comms. **1** (2010).

[11] J. Alicea, Unexpected richness in the high-field physics of graphite, Comment on [28] published in the Journal Club for Condensed Matter Physics JCCM-AUGUST-2014-01.

[12] H. Nielsen and M. Ninomiya, Phys. Lett. B **130**, 389 (1983).

[13] R. Kuechler, L. Steinke, R. Daou, M. Brando, K. Behnia, and F. Steglich, Nat. Mat. **13**, 461 (2014).

[14] K. Novoselov, A. Geim, S. Morozov, M. Katsnelson, I. Grigorieva, S. Dubonos, and A. Firsov, Nature **438**, 197 (2005).

[15] A. Castro-Neto, F. Guinea, N. Peres, K. Novoselov, and A. Geim, Rev. Mod. Phys. **81**, 109 (2009).

[16] P. R. Wallace, Phys. Rev. **71**, 622 (1946).

[17] J. W. McClure, Phys. Rev. **108**, 612 (1957).

[18] J. C. Slonczewski and P. R. Weiss, Phys. Rev. **109**, 272 (1958).

[19] J. W. McClure, Phys. Rev. **119**, 606 (1960).

[20] Y. Iye, P. M. Tedrow, G. Timp, M. Shayegan, M. S. Dresselhaus, G. Dresselhaus, A. Furukawa, and S. Tanuma, Phys. Rev. B **25**, 5478 (1982).

[21] K. Nakao, J. Phys. Soc. Jpn. **40**, 761 (1976).

[22] D. Yoshioka and H. Fukuyama, J. Phys. Soc. Japan **50**, 725 (1981).

[23] Y. Takada and H. Goto, J. Phys.: Condens. Matter **10**, 11315 (1998).

[24] H. Yaguchi and J. Singleton, Phys. Rev. Lett. **81**, 5193 (1998).

[25] H. Yaguchi and J. Singleton, J. Phys.: Condens. Matter **21**, 344207 (2009).

[26] Y. I. Latyshev, A. P. Orlov, D. Vignolles, W. Escoffier, and P. Monceau, Physica B **407**, 1885 (2012).

[27] F. Arnold, B. Yager, E. Kamper, C. Putzke, J. Nyéki, and J. Saunders, Rev. Sci. Instr. **84**, 113901 (2013).

[28] B. Fauqué, D. LeBoeuf, B. Vignolle, M. Nardone, C. Proust, and K. Behnia, Phys. Rev. Lett. **110**, 266601 (2013).

[29] K. Akiba, A. Miyake, H. Yaguchi, A. Matsuo, K. Kindo, and M. Tokunaga, J. Phys. Soc. Jpn. **84**, 054709 (2015).

[30] Z. Zhu, R. McDonald, A.Shekhter, B. Ramshaw, K. Modic, F. Balakirev, and N. Harrison, arxiv: , 1508.03645v1.

[31] Naturally Graphite, Nanotech Innovations, Michigan Technological University, 1400 Townsend Drive, Houghton, Michigan 49931-1295, United States.

[32] S. Zherlitsyn, T. Herrmannsdörfer, B. Wustmann, and J. Wosnitza, IEEE Trans. Appl. Supercon. **20**, 672 (2010).

[33] See Supplemental Material at URL for experimental and theoretical details.

[34] N. B. Brandt, G. A. Kapustin, V. G. Karavaev, A. S. Kotosonov, and E. A. Svistova, Sov. Phys.-JETP **40**, 564 (1975).

[35] K. Sugihara and J. A. Woollam, J. Phys. Soc. Jpn. **45**, 1891 (1978).

[36] H. Yaguchi, Y. Iye, T. Takamasu, and N. Miura, Physica B **184**, 332 (1993).

[37] S. Uji, J. Brooks, and Y. Iye, Physica B **246-247**, 299 (1998).

[38] We find that α, γ, α', and β', are sharp step-like transitions, whereas β is a much less well-defined transition. We thus define a more precise value for β as the extrapolated crossing field of the two linear resistive regimes either side of the transition rather than using the derivative maxima.

[39] J. Hubbard, Proc. R. Soc. Lond. A **243**, 336 (1958).

[40] K. S. Singwi, M. P. Tosi, R. H. Land, and A. Sjölander, Phys. Rev. **176**, 589 (1968).

[41] H. Yaguchi, M. Tokunaga, A. Matsuo, and K. Kindo, "Exploration of the multiple field induced density wave phase in graphite using 75-t-class pulsed magnetic fields," (2013), poster presented at the Strongly Correlated Electron Systems Conference, Aug. 5–9 2013, Tokyo, Japan.

[42] P. Lee, T. Rice, and P. Anderson, Solid State Communications **14**, 703 (1974).

[43] D. E. Moncton, J. D. Axe, and F. J. DiSalvo, Phys. Rev. Lett. **34**, 734 (1975).

[44] W. L. McMillan, Phys. Rev. B **14**, 1496 (1976).

[45] P. Bak and V. J. Emery, Phys. Rev. Lett. **36**, 978 (1976).

[46] A. D. Bruce, R. A. Cowley, and A. F. Murray, Journal of Physics C: Solid State Physics **11**, 3591 (1978).

[47] A. D. Bruce and R. A. Cowley, Journal of Physics C: Solid State Physics **11**, 3609 (1978).

[48] R. A. Cowley, Adv. Phys. **29**, 1 (1980).

[49] S. Sugai, Y. Takayanagi, and N. Hayamizu, Phys. Rev. Lett. **96**, 137003 (2006).

[50] In principle such a commensurate CDW phase should be delimited, as a function of magnetic field, by two distinct transition lines (say γ and γ') meeting each other at $T_{\rm LIT} \approx 3$ K and enclosing the value of B_γ for which the nesting vector is commensurate with the lattice. In practice, however, the window of the commensurate phase is extremely narrow and cannot be resolved experimentally, so that only a single temperature-independent transition γ can be observed (see Fig. 4).

[51] A uniform charge imbalance of 0.11 is required to explain the observed splitting, which is unphysical [29].

[52] A. H. MacDonald, Phys. Rev. Lett. **64**, 220 (1990).

[53] J. T. Chalker and A. Dohmen, Phys. Rev. Lett. **75**, 4496 (1995).

[54] L. Balents and M. P. A. Fisher, Phys. Rev. Lett. **76**, 2782 (1996).

[55] S. Cho, L. Balents, and M. P. A. Fisher, Phys. Rev. B **56**, 15814 (1997).

[56] D. P. Druist, P. J. Turley, K. D. Mara-

nowski, E. G. Gwinn, and A. C. Gossard, Phys. Rev. Lett. **80**, 365 (1998).
[57] A. F. Young, J. D. Sanchez-Yamagishi, B. HUnt, S. H. Choi, K. Watanabe, T. Taniguchi, R. C. Ashoori, and P. Jarillo-Herrero, Nature **505**, 528 (2014).
[58] M. Dzero, K. Sun, V. Galitski, and P. Coleman, Phys. Rev. Lett. **104**, 106408 (2010).
[59] M.Dzero, J. Xia, V.Galitski, and P.Coleman, Ann. Rev. Cond. Matt. Phys. **7**, 249 (2016).